

## Thermal behaviour of the system $\text{H}_3\text{BO}_3\text{--NH}_4\text{PO}_3$

E. A. Abramovich, A. F. Selevich

*Research Institute for Physical Chemical Problems, Belarusian State University, Minsk, Belarus,  
e-mail: selevich@bsu.by*

It was shown earlier that ammonium polyphosphate  $\text{NH}_4\text{PO}_3$  (APP) was a convenient reagent for preparation of normal and double ammonium-containing condensed phosphates [1]. The use of APP as phosphate-containing reagent allows to synthesize well-known compounds as well as new condensed phosphates which cannot be prepared by other techniques. In this way more than 20 well-known double ammonium-containing phosphates were synthesized in the systems  $\text{MO--NH}_4\text{PO}_3$  and  $\text{MO}_2\text{--NH}_4\text{PO}_3$ : polyphosphates  $\text{M}^{\text{II}}\text{NH}_4(\text{PO}_3)_3$  ( $\text{M}^{\text{II}} = \text{Mg, Co, Ni, Cu, Zn, Pb}$ ), cyclotriphosphates  $\text{M}^{\text{II}}\text{NH}_4\text{P}_3\text{O}_9$  ( $\text{M}^{\text{II}} = \text{Mg, Ca, Mn, Cd}$ ), polyphosphates  $\text{M}^{\text{II}}(\text{NH}_4)_2(\text{PO}_3)_4$  ( $\text{M}^{\text{II}} = \text{Mn, Co, Zn, Cd}$ ), cyclotetraphosphates  $\text{M}^{\text{II}}(\text{NH}_4)_2\text{P}_4\text{O}_{12}$  ( $\text{M}^{\text{II}} = \text{Sr, Pb}$ ), cyclooctaphosphate  $\text{Cu}_3(\text{NH}_4)_2\text{P}_8\text{O}_{24}$ , tetraphosphates  $\text{M}^{\text{IV}}(\text{NH}_4)_2\text{P}_4\text{O}_{13}$  ( $\text{M}^{\text{IV}} = \text{Si, Ge}$ ). Furthermore, 10 new compounds were also obtained in these systems. They are orthorhombic cyclotriphosphate  $\text{MnNH}_4\text{P}_3\text{O}_9$ , polyphosphates  $\text{M}^{\text{II}}(\text{NH}_4)_2(\text{PO}_3)_4$  ( $\text{M}^{\text{II}} = \text{Mg, Ca, Ni}$ ) and  $\text{Ni}(\text{NH}_4)_2(\text{PO}_3)_4 \cdot 2\text{H}_2\text{O}$ , polyphosphates  $\text{Ba}_2\text{NH}_4(\text{PO}_3)_5$  and  $\text{Ba}(\text{NH}_4)_4(\text{PO}_3)_6$ , tetraphosphates  $\text{M}^{\text{IV}}(\text{NH}_4)_2\text{P}_4\text{O}_{13}$  ( $\text{M}^{\text{IV}} = \text{Sn, Ti}$ ) [2, 3].

Double metal-ammonium condensed phosphates are of practical interest because they reveal flame retardant properties in polyamides [4, 5]. An advantage of these compounds for this application is their high thermal stability. In fact, polyamides are processed at high temperatures limiting the use of flame-retardants having low thermal stability. Unfortunately, the fire retardant properties of the majority of well-known double ammonium-containing phosphates are still deficient for real applications. Therefore, a search for new compounds of this family demonstrating higher efficiency is topical.

This work continues our systematic investigation of synthesis of trivalent metal phosphates in the systems  $\text{M}_2\text{O}_3\text{--NH}_4\text{PO}_3$  in a wide range of temperatures and reactant ratios covering the crystallization fields of phosphates of various condensation of the anion [6–8]. The aim of this study was to establish regularities of formation of boron phosphates in the APP flux, to develop a convenient technique for synthesis of detected compounds.

Thermal behavior of the  $\text{H}_3\text{BO}_3\text{--NH}_4\text{PO}_3$  system within the temperature range of 30–800 °C at the molar ratio  $\text{H}_3\text{BO}_3 : \text{NH}_4\text{PO}_3 = 1 : (1\div 10)$  using STA was studied. As a result, formation of only one boron phosphate  $\text{BPO}_4$  was determined. It was shown by XRD that this compound was a tetragonal polymorph of  $\text{BPO}_4$  [9]. No double ammonium-containing boron phosphates were detected. Finally, the “poorness” of boron phosphate chemistry is considered.

### References

1. A. F. Selevich, G. F. Levchik, A. I. Lesnikovich, S. V. Levchik. Belarus. Pat. Appl. (1994) 260A.
2. A. F. Selevich, L. S. Ivashkevich, K. I. Khurs, A. S. Lyakhov, A. I. Lesnikovich. Phosphorus Res. Bull. (2002) 13: 227.
3. A. F. Selevich, E. V. Grushevich, L. S. Ivashkevich, G. L. Krul, N. V. Ilushchenko, A. I. Lesnikovich. Phosphorus Res. Bull. (2005) 19: 228.
4. S. Duquesne, M. Le Bras, S. Bourbigot, R. Delobel. Phosphorus Res. Bull. (1999) 10: 88.
5. A. I. Balabanovich, A. I. Lesnikovich, A. F. Selevich. Belarus. Pat. Appl. a20120476. 2012.
6. E. A. Abramovich, A. I. Lesnikovich, A. F. Selevich. Sviridov readings (2009) 5: 67.
7. E. A. Abramovich, A. I. Lesnikovich, A. F. Selevich. Sviridov readings (2011) 7: 9.
8. E. A. Abramovich, A. I. Lesnikovich, A. F. Selevich. Sviridov readings. (2013) 9: 11.
9. G. E. R. Schulze. Z. Phys. Chem. (1934) 24 (2): 215.